ELSEVIER

Journal of Immunological Methods 182 (1995) 263–264

JOURNAL OF IMMUNOLOGICAL METHODS

Contents to volume 182 (1995)

Number 1 – 15 May

Research reports

Re-evaluation of the storage conditions for blood samples which are used for determination of complement activation
S. Stöve, A. Klos, W. Bautsch and J. Köhl (Germany) 1

Phage display used for gene cloning of human recombinant antibody against the erythrocyte surface antigen, rhesus D
M. Dziegiel, L.K. Nielsen, P.S. Andersen, A. Blancher, E. Dickmeiss and J. Engberg (Denmark, France) 7

Establishment of a T cell-dependent nude mouse liver injury model induced by *Propionibacterium acnes* and LPS
Y. Tanaka, A. Takahashi, K. Kobayashi, I. Arai, S. Higuchi, S. Otomo, K. Watanabe, S. Habu and T. Nishimura (Japan) 21

Characterization of the interaction of alkaline phosphatase with an activity inhibiting monoclonal antibody by progress curve analysis
G.A. Cumme, U. Walter, R. Bublitz, H. Hoppe, H. Rhode and A. Horn (Germany) 29

Phage display of disulfide-stabilized Fv fragments
U. Brinkmann, P.S. Chowdhury, D.M. Roscoe and I. Pastan (USA) 41

Pristane induces high titers of anti-Su and anti-nRNP/Sm autoantibodies in BALB/c mice Quantitation by antigen capture ELISAs based on monospecific human autoimmune sera
M. Satoh, E.L. Treadwell and W.H. Reeves (USA) 51

Antigenicity of cystatin-binding proteins from parasitic protozoan. Detection by a proteinase inhibitor based capture immunoassay (PINC-ELISA)
J. Scharfstein, M. Abrahamson, C.B. Palatnik de Souza, A. Barral and I.V. Silva (Brazil, Sweden) 63

Use of the Pannell-Milstein roller bottle apparatus to produce high concentrations of the CSF-1, the mouse macrophage growth factor
E. Olivas, B.B.D.-M. Chen and W.S. Walker (USA) 73

Development of an enzyme immunoassay for rat soluble interleukin-2 receptors
A.G. Pockley, S. Williams, S.D. Reid and M.J. Bowles (UK) 81

Analysis of anticholesterol antibodies using hydrophobic membranes
J. Aniagolu, G.M. Swartz, Jr., J. Dijkstra, J.W. Madsen, J.J. Raney and S.J. Green (USA) 85

In situ hybridization for cytokine mRNA with digoxigenin-labeled riboprobes. Sensitivity of detection and double label applications
L.J. Karr, A. Panoskaltsis-Mortari, J. Li, D. Devore-Carter, C.T. Weaver and R.P. Bucy (USA) 93

Assaying tumor necrosis factor concentrations in human serum A WHO International Collaborative Study
S. De Kossodo, V. Houba, G.E. Grau and WHO Collaborative Study Group (USA, Switzerland) 107

Purification of human basophils and mast cells by multistep separation technique and mAb to CDw17 and CD117/c-kit
M. Willheim, H. Agis, W.R. Sperr, M. Köller, H.-C. Bankl, H. Kiener, G. Fritsch, W. Füreder, A. Spittler, W. Graninger, O. Scheiner, H. Gadner, K. Lechner, G. Boltz-Nitulescu and P. Valent (Austria) 115

Quantification and mapping of antigenic determinants of serum amyloid A (SAA) protein utilizing sequence-specific immunoglobulins and Eu^{3+} as a specific probe for time-resolved fluorometric immunoassay
E. Malle, G. Münscher, T. Müller, H. Vermeer and A. Ibovnik (Austria, Germany) 131

Letters to the editors

Under proper experimental conditions the solid-phase antigen does not disrupt the liquid phase equilibrium when measuring dissociation constants by competition ELISA
B. Friguet, A.F. Chaffotte, L. Djavadi-Ohaniance and M.E. Goldberg (France) 145

Reply to the letter of B. Friguet et al.
S.J. Seligman (USA) 149

Announcement 151

Notes to authors 153

Number 2 – 9 June 1995

Research reports
Secretion capture and report web: use of affinity derivatized agarose microdroplets for the selection of hybridoma cells
F. Gray, J.S. Kenney and J.F. Dunne (USA) 155
A novel method to purify immunotoxins from free antibodies using modified recombinant toxins
M. Dhanabal, D.K. Fryxell and S. Ramakrishnan (USA) 165
In vitro proliferative responses of human peripheral blood mononuclear cells to non-recall antigens
J.L. Young, A. Daser and P.C.L. Beverley (UK) 177
Elution of human IgG from affinity columns containing immobilised variants of protein A
S.P. Bottomley, B.J. Sutton and M.G. Gore (UK) 185
Simultaneous three-color analysis of the surface phenotype and DNA-RNA quantitation using 7-amino-actinomycin D and pyronin Y
K. Toba, E.F. Winton, T. Koike and A. Shibata (Japan, USA) 193
Efficient isolation of mutant antigen presenting cell lines by functional selection using T cell clones
S. Koarada, E. Kubota, M. Tokushima, K. Naitoh, K. Miyake and M. Kimoto (Japan) 209
Monopalmitic acid-peptide conjugates induce cytotoxic T cell responses against malarial epitopes: importance of spacer amino acids
A.F.M. Verheul, V. Udhayakumar, D.L. Jue, R.M. Wohlhueter and A.A. Lal (USA) 219
Grafting peptides onto polystyrene microplates for ELISA
A. Niveleau, C. Bruno, E. Drouet, R. Brebant, A. Sergeant and F. Troalen (France) 227
The derivation of binding parameters from effector and target conjugate frequency data using linear and non-linear data-fitting transformations. Application of such transformations to the NK-MOLT4 and NK-K562 effector-target systems
P. Garcia-Peñarrubia, L. Cabrera, F. Lajarin and J. Galvez (Spain) 235
Cytokine immunotrapping: an assay to study the kinetics of production and consumption or degradation of human interferon-γ
A.C. Akdiş, H. Towbin, P. Libsig, J. Motz and Ş.Ş. Alkan (Switzerland) 251

Contents to volume 182 (1995) 263

Notes to authors 265